

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mouse anti Human CD14, conjugated with FITC, IgG2b, Clone: [FWKW], Monoclonal NMB-0142

Artikelname	Mouse anti Human CD14, conjugated with FITC, IgG2b, Clone: [FWKW], Monoclonal
Artikelnummer	NMB-0142
Hersteller Artikelnummer	0142
Alternativnummer	NMB-0142
Hersteller	NordicMubio
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC
Spezies Reaktivität	Human
Konjugation	FITC
Produktbeschreibung	Identification of human Monocyte/Macrophage cells expressing the 55 kDa M.W. surface antigen. CD14 is present on 70-93% of Monocyte/Macrophage cells in normal peripheral blood....
Klonalität	Monoclonal
Konzentration	Titered for flow cytometry
Klon-Bezeichnung	[FWKW]
Isotyp	IgG2b

UniProt	P08571
Puffer	Provided as solution in phosphate buffered saline with 0.08% sodium azide and 0.2% carrier protein
Reinheit	Protein A/G Chromatography
Formulierung	FITC
Formel	Provided as solution in phosphate buffered saline with 0.08% sodium azide and 0.2% carrier protein
Anwendungsbeschreibung	PBMC: Add 10 µl of MAB/10⁶ PBMC in 100 µl PBS. Mix gently and incubate for 15 minutes at 2 to 8°C. Wash twice with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. WHOLE BLOOD: Add 10 µl of MAB/100 µl of whole blood. Mix gently and incubate for 15 minutes at room temperature (20°C). Lyse the whole blood. Wash once with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. See instrument manufacturers instructions for Lysed Whole Blood and Immunofluorescence analysis with a flow cytometer or microscope. ALLOPHYCOCYANIN: (APC) conjugates are analyzed in multi-color flow cytometry with instruments equipped with a second laser and proper filters. Laser excitation is at 633 nm with a Helium Neon (HeNe) laser or a 600-640 nm (633 nm) range for a Dye laser. Peak fluorescence emission is at 660 nm.