

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mouse anti Human CD25, conjugated with FITC, IgG1, Clone: [ITYV], Monoclonal NMB-0252

| | |
|--------------------------|--|
| Artikelname | Mouse anti Human CD25, conjugated with FITC, IgG1, Clone: [ITYV], Monoclonal |
| Artikelnummer | NMB-0252 |
| Hersteller Artikelnummer | 0252 |
| Alternativnummer | NMB-0252 |
| Hersteller | NordicMubio |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC |
| Spezies Reaktivität | Human |
| Immunogen | CD25=Derived from the hybridization of mouse NS-1 myeloma cells with spleen cells of BALB/c mice immunized with PHA-activated human T-lymphocytes. |
| Konjugation | FITC |
| Produktbeschreibung | Identification of human receptor for Interleukin-2 (IL-2R) expressing the 55,000 M.W. surface antigen.... |
| Klonalität | Monoclonal |
| Konzentration | Titered for flow cytometry |
| Klon-Bezeichnung | [ITYV] |

| | |
|------------------------|--|
| Isotyp | IgG1 |
| UniProt | P01589 |
| Puffer | Provided as solution in phosphate buffered saline with 0.08% sodium azide and 0.2% carrier protein |
| Reinheit | Protein A/G Chromatography |
| Formulierung | FITC |
| Formel | Provided as solution in phosphate buffered saline with 0.08% sodium azide and 0.2% carrier protein |
| Anwendungsbeschreibung | <p>PBMC: Add 10 µl of MAB/10 6 PBMC in 100 µl PBS. Mix gently and incubate for 15 minutes at 2 to 8C. Wash twice with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. WHOLE BLOOD: Add 10 µl of MAB/100 µl of whole blood. Mix gently and incubate for 15 minutes at room temperature 20C. Lyse the whole blood. Wash once with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. See instrument manufacturers instructions for Lysed Whole Blood and Immunofluorescence analysis with a flow cytometer or microscope.</p> |