

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mouse anti Human Keratin (4,5,6,8,10,13,18), IgG1, Clone: [C11], Unconjugated, Monoclonal NMB-X1260M

| | |
|--------------------------|---|
| Artikelname | Mouse anti Human Keratin (4,5,6,8,10,13,18), IgG1, Clone: [C11], Unconjugated, Monoclonal |
| Artikelnummer | NMB-X1260M |
| Hersteller Artikelnummer | X1260M |
| Alternativnummer | NMB-X1260M |
| Hersteller | NordicMubio |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Konjugation | Unconjugated |
| Produktbeschreibung | Cytokeratins (CK) are intermediate filaments of epithelial cells, both in keratinizing tissue (ie., skin) and non-keratinizing cells (ie., mesothelial cells). Although not a traditional marker for endothelial cells, cytokeratins have also been found ... |
| Klonalität | Monoclonal |
| Konzentration | See vial for concentration |
| Klon-Bezeichnung | [C11] |
| Isotyp | IgG1 |

| | |
|------------------------|---|
| UniProt | P13647 |
| Puffer | Provided as solution in phosphate buffered saline with 0.08% sodium azide |
| Quelle | Hybridoma produced by the fusion of splenocytes from mice immunized with cytoskeleton preparation from human A431 carcinoma cells and mouse myeloma cells. |
| Reinheit | Purified by saturated ammonium sulphate precipitation |
| Formulierung | Unconjugated |
| Formel | Provided as solution in phosphate buffered saline with 0.08% sodium azide |
| Anwendungsbeschreibung | X1260M detects keratins 4,5,6,8,10,13 & 18 by Western blot. Immunohistochemistry can be performed on frozen tissues and paraffin sections. Optimal concentration should be evaluated by serial dilutions. X1260M reacts with a variety of normal and neoplastic epithelia. Reacting with simple epithelium and both basal and superbasal layers of cornifying and non-cornifying squamous epithelium this antibody is also useful in staining cultured epithelial cell lines. It is useful in differentiating epithelial tumors from non-epithelial tumors. |