

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mouse anti Lamin A, IgG3, Clone: [133A2], Monoclonal NMB-MUB1101P

Artikelname	Mouse anti Lamin A, IgG3, Clone: [133A2], Monoclonal
Artikelnummer	NMB-MUB1101P
Hersteller Artikelnummer	MUB1101P
Alternativnummer	NMB-MUB1101P
Hersteller	NordicMubio
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, ICC, IHC-Fr, IHC-P, WB
Spezies Reaktivität	Bovine, Canine, Human, Mouse, Rat
Produktbeschreibung	Nuclear lamins form a network of intermediate-type filaments at the nucleoplasmic site of the nuclear membrane. Two main subtypes of nuclear lamins can be distinguished, i.e. A-type lamins and B-type lamins. The A-type lamins comprise a set of three ...
Klonalität	Monoclonal
Klon-Bezeichnung	[133A2]
Isotyp	IgG3
UniProt	P02545

Puffer	Each vial contains 100 ul 1 mg/ml purified monoclonal antibody in PBS containing 0.09% sodium azide. Since this IgG3 antibody has to be purified from the culture medium by ion-exchange chromatography it will contain remnants of bovine serum albumin.
Quelle	133A2 is a mouse monoclonal IgG3/kappa antibody obtained from fusion of P3/X63.Ag8.653 mouse myeloma cells with spleen cells from a BALB/c mouse immunized with partially purified recombinant human lamin A.
Formel	Each vial contains 100 ul 1 mg/ml purified monoclonal antibody in PBS containing 0.09% sodium azide. Since this IgG3 antibody has to be purified from the culture medium by ion-exchange chromatography it will contain remnants of bovine serum albumin.
Anwendungsbeschreibung	133A2 is suitable for immunoblotting, immunocytochemistry on permeabilized cells, immunohistochemistry on frozen sections and paraffin embedded tissues, and flow cytometry. Optimal antibody dilution should be determined by titration, recommended range is 1:100 - 1:200 for immunocytochemistry, and immunohistochemistry with fluorescent secondary antibodies or avidin-biotinylated horseradish peroxidase complex (ABC) as detection reagent, and 1:100 - 1:1000 for immunoblotting applications.