

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Rabbit anti Human Phosphatidic Acid Phosphatase Domain Containing 2 (PPAPDC2), Unconjugated NMB-X1717P

|                          |   |
|--------------------------|---|
| Artikelname              | Rabbit anti Human Phosphatidic Acid Phosphatase Domain Containing 2 (PPAPDC2), Unconjugated   |
| Artikelnummer            | NMB-X1717P  |
| Hersteller Artikelnummer | X1717P  |
| Alternativnummer         | NMB-X1717P  |
| Hersteller               | NordicMubio   |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | IP, WB  |
| Spezies Reaktivität      | Human   |
| Immunogen                | Synthetic peptide derived from human phosphatidic acid phosphatase type 2 domain containing 2 (PPAPDC2)   |
| Konjugation              | Unconjugated  |
| Produktbeschreibung      | Upon cell activation Presqualene diphosphate (PSDP)2, a bioactive lipid, rapidly remodels to presqualene monophosphate (PSMP). A newly identified and characterized phosphatase (phosphatidic acid phosphatase domain containing 2 (PPAPDC2)) has recent,y... |
| Konzentration            | Lot specific, see vial  |
| UniProt                  | <a href="#">Q8IY26</a>  |

|                        |  |
|------------------------|--|
| Puffer                 | Provided as ligand affinity purified antibody in phosphate buffered saline with 0.1% BSA and 0.08% sodium azide                  |
| Formulierung           | Unconjugated   |
| Formel                 | Provided as ligand affinity purified antibody in phosphate buffered saline with 0.1% BSA and 0.08% sodium azide                  |
| Anwendungsbeschreibung | Antibody can be used for Western blotting (1:400-1:800 dilution). Optimal concentration should be evaluated by serial dilutions. |