

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD28 Antibody, Clone: [C28/75], Mouse, Monoclonal NSJ-V2053SAF-100UG

Artikelname	CD28 Antibody, Clone: [C28/75], Mouse, Monoclonal
Artikelnummer	NSJ-V2053SAF-100UG
Hersteller Artikelnummer	V2053SAF-100UG
Alternativnummer	NSJ-V2053SAF-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	FACS, Functional Studies
Spezies Reaktivität	Human
Immunogen	Recombinant human CD28 protein was used as the immunogen for this antibody.
Produktbeschreibung	This antibody recognizes a glycoprotein of 44-88kDa, which is identified as CD28. It is the critical T-cell co-stimulatory receptor which provides to the cell the important second activation signal by binding CD80 and CD86 that are expressed by antig...
Klonalität	Monoclonal
Konzentration	1 mg/ml in 1X PBS, BSA free, sodium azide free
Klon-Bezeichnung	[C28/75]
Isotyp	Mouse IgG1, kappa

NCBI	940
Puffer	1 mg/ml in 1X PBS, BSA free, sodium azide free
Reinheit	Protein G affinity chromatography
Formulierung	1 mg/ml in 1X PBS, BSA free, sodium azide free
Target-Kategorie	CD28
Antibody Type	Primary Antibody
Application Verdünnung	Flow cytometry: 1-2ug/10 ⁶ cells, Functional studies (order BSA/sodium azide-free format)
Anwendungsbeschreibung	The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the antibody to be titrated up or down for optimal performance. 1. Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 1mM EDTA Buffer, pH 8.5-9.5, for 10-20 min followed by cooling at RT for 20 minutes.