

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

c-Myc Antibody, Clone: [MVH39-1], Mouse, Monoclonal NSJ-V7021IHC-7ML

Artikelname	c-Myc Antibody, Clone: [MVH39-1], Mouse, Monoclonal
Artikelnummer	NSJ-V7021IHC-7ML
Hersteller Artikelnummer	V7021IHC-7ML
Alternativnummer	NSJ-V7021IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	A C-terminal peptide was used as the immunogen for the c-Myc antibody.
Produktbeschreibung	Transcription factor that binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5-CAC[GA]TG-3. Activates the transcription of growth-related genes. [UniProt]...
Klonalität	Monoclonal
Konzentration	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide, *For IHC use only*
Klon-Bezeichnung	[MVH39-1]
Isotyp	Mouse IgG1, kappa

UniProt	P01106
Puffer	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide, *For IHC use only*
Reinheit	Protein G affinity chromatography
Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Target-Kategorie	c-Myc
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Titering of the c-Myc antibody may be required for optimal performance.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.