

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Glypican-3 Antibody, Clone: [SGPN3-1], Mouse, Monoclonal NSJ-V7027-20UG

Artikelname	Glypican-3 Antibody, Clone: [SGPN3-1], Mouse, Monoclonal
Artikelnummer	NSJ-V7027-20UG
Hersteller Artikelnummer	V7027-20UG
Alternativnummer	NSJ-V7027-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	FACS, IF, IHC-P
Spezies Reaktivität	Human, Rat
Immunogen	Recombinant full-length human GPC3 protein was used as the immunogen for the Glypican-3 antibody.
Produktbeschreibung	Glypican-3 is a proteoglycan that bears heparan sulfate. Inhibits the dipeptidyl peptidase activity of DPP4. May be involved in the suppression/modulation of growth in the predominantly mesodermal tissues and organs. May play a role in the modulation...
Klonalität	Monoclonal
Konzentration	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Klon-Bezeichnung	[SGPN3-1]
Isotyp	Mouse IgG1, kappa

UniProt	P51654
Puffer	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Reinheit	Protein G affinity chromatography
Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Target-Kategorie	Glypican 3
Antibody Type	Primary Antibody
Application Verdünnung	Flow cytometry: 1-2ug/million cells,Immunofluorescence: 1-2ug/ml,Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT (1),Prediluted IHC only format: incubate for 30 min at RT (2)
Anwendungsbeschreibung	Optimal dilution of the Glypican-3 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.