

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

AMPD3 Antibody, Clone: [ETAD3-1], Mouse, Monoclonal NSJ-V7030-20UG

Artikelname	AMPD3 Antibody, Clone: [ETAD3-1], Mouse, Monoclonal
Artikelnummer	NSJ-V7030-20UG
Hersteller Artikelnummer	V7030-20UG
Alternativnummer	NSJ-V7030-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human protein was used as the immunogen for the AMPD3 antibody.
Produktbeschreibung	AMPD3 is a member of the AMP deaminase gene family. It is a highly regulated enzyme that catalyzes the hydrolytic deamination of adenosine monophosphate to inosine monophosphate, a branch point in the adenylate catabolic pathway. This gene encodes th...
Klonalität	Monoclonal
Konzentration	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Klon-Bezeichnung	[ETAD3-1]
Isotyp	Mouse IgG2b, kappa

UniProt	Q01432
Puffer	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Reinheit	Protein G affinity chromatography
Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Target-Kategorie	AMPD3
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistology (FFPE): 2-4ug/ml for 30 min at RT (1),Prediluted IHC only format: incubate for 30 min at RT (2)
Anwendungsbeschreibung	Optimal dilution of the AMPD3 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.