

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Blood Group Antigen A (CD173) Antibody, IgM, Clone: [HE-14], Mouse, Monoclonal NBT-28-MSM4-P2

| | |
|--------------------------|---|
| Artikelname | Blood Group Antigen A (CD173) Antibody, IgM, Clone: [HE-14], Mouse, Monoclonal |
| Artikelnummer | NBT-28-MSM4-P2 |
| Hersteller Artikelnummer | 28-MSM4-P2 |
| Alternativnummer | NBT-28-MSM4-P2-100,NBT-28-MSM4-P2-200 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IF |
| Spezies Reaktivität | Human |
| Immunogen | Mixture of erythrocytes of blood group A1and glycoprotein fraction isolated from the saliva of secretors with blood group A |
| Produktbeschreibung | This antibody recognizes human blood group A (monofucosyl and difucosyl A antigens with chain types 1, 2, 3, 4, 5, 6) and Forssmann antigen. It is also reactive with the immuno-dominant A trisaccharide. Blood group antigen expression in human colon c... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [HE-14] |
| Molekulargewicht | Multiple |

| | |
|------------------------|--|
| Isotyp | IgM |
| NCBI | 28 |
| UniProt | P16442 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Agglutination, Immunofluorescence (2-4ug/ml), Immunohistology (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for |