

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant CD25 / IL2RA (Activated Lymphocyte Marker) Antibody, Clone: [IL2RA/4375R], Rabbit, Monoclonal NBT-3559-RBM9-P1ABX

| | |
|--------------------------|---|
| Artikelname | Recombinant CD25 / IL2RA (Activated Lymphocyte Marker) Antibody, Clone: [IL2RA/4375R], Rabbit, Monoclonal |
| Artikelnummer | NBT-3559-RBM9-P1ABX |
| Hersteller Artikelnummer | 3559-RBM9-P1ABX |
| Alternativnummer | NBT-3559-RBM9-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment (around aa1-250) corresponding to human IL-2 receptor alpha |
| Produktbeschreibung | CD25, or interleukin 2 (IL2) receptor alpha (IL2RA), and beta (IL2RB) chains, together with the common gamma chain (IL2RG), constitute the high-affinity IL2 receptor. Homodimeric alpha chains (CD25) result in low-affinity receptor, while homodimeric ... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [IL2RA/4375R] |
| Molekulargewicht | 31kDa |

| | |
|------------------------|--|
| NCBI | 3559 |
| UniProt | P01589 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes),Optimal dilutio |