

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 13 (Non-Keratinized Squamous Epithelial Marker) Antibody, IgG1, Clone: [KRT13/2850], Mouse, Monoclonal NBT-3860-MSM4-P0

| | |
|--------------------------|--|
| Artikelname | Cytokeratin 13 (Non-Keratinized Squamous Epithelial Marker) Antibody, IgG1, Clone: [KRT13/2850], Mouse, Monoclonal |
| Artikelnummer | NBT-3860-MSM4-P0 |
| Hersteller Artikelnummer | 3860-MSM4-P0 |
| Alternativnummer | NBT-3860-MSM4-P0-20,NBT-3860-MSM4-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IF, IHC, WB |
| Spezies Reaktivität | Human, Mouse, Rabbit, Rat |
| Immunogen | esophageal keratins of rabbit origin |
| Produktbeschreibung | Cytokeratins comprise a diverse group of intermediate filament proteins (IFPs) that are expressed as pairs in both keratinized and non-keratinized epithelial tissue. Cytokeratins play a critical role in differentiation and tissue specialization and fu... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [KRT13/2850] |
| Molekulargewicht | 52kDa |
| Isotyp | IgG1 |

| | |
|------------------------|--|
| NCBI | 3860 |
| UniProt | P13646 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes),Optimal dilutio |