

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Recombinant MMP3 (Marker of Metastasis and Rheumatoid Arthritis) Antibody, IgG1, Clone: [rMMP3/1730], Mouse, Monoclonal NBT-4314-MSM4-P1ABX**

|                          |   |
|--------------------------|---|
| Artikelname              | Recombinant MMP3 (Marker of Metastasis and Rheumatoid Arthritis) Antibody, IgG1, Clone: [rMMP3/1730], Mouse, Monoclonal   |
| Artikelnummer            | NBT-4314-MSM4-P1ABX   |
| Hersteller Artikelnummer | 4314-MSM4-P1ABX   |
| Alternativnummer         | NBT-4314-MSM4-P1ABX-100   |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant full-length human MMP3 protein  |
| Produktbeschreibung      | The matrix metalloproteinases (MMP) are a family of peptidase enzymes responsible for the degradation of extracellular matrix components, including collagen, gelatin, fibronectin, laminin and proteoglycan. Transcription of MMP genes is differentially... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [rMMP3/1730]  |
| Molekulargewicht         | 57kDa   |
| Isotyp                   | IgG1  |

|                        |   |
|------------------------|---|
| NCBI                   | <a href="#">4314</a>  |
| UniProt                | <a href="#">P08254</a>  |
| Formulierung           | 200ug/ml of Ab Purified by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.  |
| Antibody Type          | Recombinant Monoclonal Antibody   |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution f |