

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **RNA Polymerase II CTD Repeat YSPTSPS (Phospho S5) Antibody, IgM, Clone: [CTD 8A7], Mouse, Monoclonal NBT-5430-MSM2-P1BX**

|                          |   |
|--------------------------|---|
| Artikelname              | RNA Polymerase II CTD Repeat YSPTSPS (Phospho S5) Antibody, IgM, Clone: [CTD 8A7], Mouse, Monoclonal  |
| Artikelnummer            | NBT-5430-MSM2-P1BX  |
| Hersteller Artikelnummer | 5430-MSM2-P1BX  |
| Alternativnummer         | NBT-5430-MSM2-P1BX-100  |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FC, IF, IHC, WB   |
| Spezies Reaktivität      | Human, Yeast  |
| Immunogen                | Ten repeats of synthetic peptide YSPTSPS using chemically synthesized phospho-Ser5  |
| Produktbeschreibung      | RNA polymerase II (Pol II) is an enzyme that is composed of 12 subunits and is responsible for the transcription of protein-coding genes. Transcription initiation requires Pol II-mediated recruitment of transcription machinery to a target promoter, t... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [CTD 8A7]   |
| Molekulargewicht         | 192-253kDa  |

|                        |  |
|------------------------|--|
| Isotyp                 | IgM  |
| NCBI                   | <a href="#">5430</a>   |
| UniProt                | <a href="#">P24928</a>   |
| Formulierung           | 200ug/ml of Ab Purified from Bioreactor Concentrate. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.  |
| Antibody Type          | Monoclonal Antibody  |
| Anwendungsbeschreibung | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA |