

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Calcineurin B / PPP3R1 Antibody, IgG1, Clone: [CALNB/2342], Mouse, Monoclonal NBT-5534-MSM2-P1ABX

| | |
|--------------------------|---|
| Artikelname | Calcineurin B / PPP3R1 Antibody, IgG1, Clone: [CALNB/2342], Mouse, Monoclonal |
| Artikelnummer | NBT-5534-MSM2-P1ABX |
| Hersteller Artikelnummer | 5534-MSM2-P1ABX |
| Alternativnummer | NBT-5534-MSM2-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length Calcineurin B protein |
| Produktbeschreibung | Calcineurin is an enzyme that dephosphorylates serine and threonine residues in proteins. It is a heterodimer of a 59kDa catalytic A subunit and a 19kDa regulatory B subunit that is activated by the binding of calcium ions and calmodulin. Calcineurin... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [CALNB/2342] |
| Molekulargewicht | 19kDa |

| | |
|------------------------|---|
| Isotyp | IgG1 |
| NCBI | 5534 |
| UniProt | P63098 |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution for a |