

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **KLK7 (Kallikrein Related Peptidase 7) Antibody, IgG2b, Clone: [KLK7/4691], Mouse, Monoclonal NBT-5650-MSM1-P0**

|                          |   |
|--------------------------|---|
| Artikelname              | KLK7 (Kallikrein Related Peptidase 7) Antibody, IgG2b, Clone: [KLK7/4691], Mouse, Monoclonal  |
| Artikelnummer            | NBT-5650-MSM1-P0  |
| Hersteller Artikelnummer | 5650-MSM1-P0  |
| Alternativnummer         | NBT-5650-MSM1-P0-20,NBT-5650-MSM1-P0-100  |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant fragment of human KLK7 protein (around aa 1-200) (exact sequence is proprietary)  |
| Produktbeschreibung      | Kallikreins (KLKs) belong to the serine protease family of proteolytic enzymes. Human pancreatic/renal KLK encodes for the KLK1 enzyme, which is involved in post-translational processing of polypeptide precursors. The function of the other members of... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [KLK7/4691]   |
| Molekulargewicht         | 27kDa   |

|                        |  |
|------------------------|--|
| Isotyp                 | IgG2b  |
| NCBI                   | <a href="#">5650</a>   |
| UniProt                | <a href="#">P49862</a>   |
| Formulierung           | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.   |
| Antibody Type          | Monoclonal Antibody  |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes),Optimal dilutio |