

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

STAT5B Antibody, IgG1, Clone: [STAT5B/2611], Mouse, Monoclonal NBT-6777-MSM1-P1ABX

| | |
|--------------------------|---|
| Artikelname | STAT5B Antibody, IgG1, Clone: [STAT5B/2611], Mouse, Monoclonal |
| Artikelnummer | NBT-6777-MSM1-P1ABX |
| Hersteller Artikelnummer | 6777-MSM1-P1ABX |
| Alternativnummer | NBT-6777-MSM1-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human STAT5B protein |
| Produktbeschreibung | Signal transducer and activator of transcription 5A (Stat5a) and Stat5b, which share 96% homology, undergo receptor tyrosine kinase or G protein-coupled receptor-dependent phosphorylation in response to cytokines or growth factors, and then form homo... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [STAT5B/2611] |
| Molekulargewicht | 94kDa |
| Isotyp | IgG1 |
| NCBI | 6777 |

| | |
|------------------------|---|
| UniProt | P51692 |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution for a |