

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker) Antibody, Clone: [VWF/7979R], Rabbit, Monoclonal NBT-7450-RBM20-P1ABX

| | |
|--------------------------|---|
| Artikelname | Recombinant von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker) Antibody, Clone: [VWF/7979R], Rabbit, Monoclonal |
| Artikelnummer | NBT-7450-RBM20-P1ABX |
| Hersteller Artikelnummer | 7450-RBM20-P1ABX |
| Alternativnummer | NBT-7450-RBM20-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment (around aa 1150-1250) of human vWF protein (exact sequence is proprietary) |
| Produktbeschreibung | Human von Willebrand factor (VWF or factor VIII-related antigen) is a 309 kDa multimeric plasma glycoprotein, important in the maintenance of hemostasis, it promotes adhesion of platelets to the sites of vascular injury by forming a molecular bridge ... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [VWF/7979R] |
| Molekulargewicht | 250kDa |

| | |
|------------------------|--|
| NCBI | 7450 |
| UniProt | P04275 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes),Optimal dilutio |