

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **XRCC3 Antibody, IgG1, Clone: [10F1/6], Mouse, Monoclonal NBT-7517-MSM1-P1ABX**

|                          |   |
|--------------------------|---|
| Artikelname              | XRCC3 Antibody, IgG1, Clone: [10F1/6], Mouse, Monoclonal  |
| Artikelnummer            | NBT-7517-MSM1-P1ABX   |
| Hersteller Artikelnummer | 7517-MSM1-P1ABX   |
| Alternativnummer         | NBT-7517-MSM1-P1ABX-100   |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC, WB   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant full-length human XRCC3 protein   |
| Produktbeschreibung      | The x-ray repair cross-complementing (XRCC) proteins are responsible for efficiently repairing and maintaining genetic stability following DNA base damage. These genes share sequence similarity with the yeast DNA repair protein Rad51. XRCC1 is a prot... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [10F1/6]  |
| Molekulargewicht         | 40kDa   |
| Isotyp                   | IgG1  |
| NCBI                     | <a href="#">7517</a>  |

|                        |   |
|------------------------|---|
| UniProt                | <a href="#">O43542</a>  |
| Formulierung           | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.  |
| Antibody Type          | Monoclonal Antibody   |
| Anwendungsbeschreibung | Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes |