

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD2 / Lymphocyte Function Antigen 2 (LFA-2) Antibody, IgG2b, Clone: [HuLy-m1], Mouse, Monoclonal NBT-914-MSM4-P0**

|                          |   |
|--------------------------|---|
| Artikelname              | CD2 / Lymphocyte Function Antigen 2 (LFA-2) Antibody, IgG2b, Clone: [HuLy-m1], Mouse, Monoclonal  |
| Artikelnummer            | NBT-914-MSM4-P0   |
| Hersteller Artikelnummer | 914-MSM4-P0   |
| Alternativnummer         | NBT-914-MSM4-P0-20,NBT-914-MSM4-P0-100  |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FC, Functional Studies, IF  |
| Spezies Reaktivität      | Feline, Human   |
| Immunogen                | Human thymocytes  |
| Produktbeschreibung      | CD2 interacts through its amino-terminal domain with the extracellular domain of CD58 (also designated CD2 ligand) to mediate cell adhesion. CD2/CD58 binding can enhance antigen-specific T cell activation. CD2 is a transmembrane glycoprotein that is ... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [HuLy-m1]   |
| Molekulargewicht         | ~50kDa  |

|                        |  |
|------------------------|--|
| Isotyp                 | IgG2b  |
| NCBI                   | <a href="#">914</a>  |
| UniProt                | <a href="#">P06729</a>   |
| Formulierung           | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.                 |
| Antibody Type          | Monoclonal Antibody  |
| Anwendungsbeschreibung | Functional Studies (Order Ab without Azide), ,Flow Cytometry (1-2ug/million cells), ,Immunofluorescence (1-2ug/ml), ,Optimal dilution for a specific application should be determined. |