

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant CD27 (Tumor Necrosis Factor Receptor Superfamily 7) Antibody, Clone: [LPFS2/2034R], Rabbit, Monoclonal NBT-939-RBM4-P0

| | |
|--------------------------|---|
| Artikelname | Recombinant CD27 (Tumor Necrosis Factor Receptor Superfamily 7) Antibody, Clone: [LPFS2/2034R], Rabbit, Monoclonal |
| Artikelnummer | NBT-939-RBM4-P0 |
| Hersteller Artikelnummer | 939-RBM4-P0 |
| Alternativnummer | NBT-939-RBM4-P0-20,NBT-939-RBM4-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human full-length CD22 protein (exact sequence is proprietary) |
| Produktbeschreibung | Recognizes a protein of a disulfide-linked 120kDa dimer, identified as CD27. It is expressed on the majority of peripheral T cells, medullary thymocytes, memory-type B cells, and natural killer cells. It is a transmembrane phosphoglycoprotein that be... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [LPFS2/2034R] |
| Molekulargewicht | 120kDa |

| | |
|------------------------|--|
| NCBI | 939 |
| UniProt | P26842 |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris buffer with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes)Optimal di |