

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD30 / TNFRSF8 (Hodgkin & Reed-Sternberg Cell Marker) Antibody, IgG1, Clone: [Ki-1/779], Mouse, Monoclonal NBT-943-MSM3-P1ABX

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | CD30 / TNFRSF8 (Hodgkin & Reed-Sternberg Cell Marker) Antibody, IgG1, Clone: [Ki-1/779], Mouse, Monoclonal |
| Artikelnummer | NBT-943-MSM3-P1ABX |
| Hersteller Artikelnummer | 943-MSM3-P1ABX |
| Alternativnummer | NBT-943-MSM3-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human TNFRSF8 protein |
| Produktbeschreibung | Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease,... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [Ki-1/779] |
| Molekulargewicht | 105-120kDa |

| | |
|------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Isotyp | IgG1 |
| NCBI | 943 |
| UniProt | P28908 |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 1mM EDTA, pH 7.5-8.5, for 45 min at 95degC |