

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD40 Ligand / CD154 / TRAP1 (Activation Marker of T-Lymphocytes) Antibody, IgG2b, Clone: [CD40LG/2761], Mouse, Monoclonal NBT-959-MSM1-B1

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | CD40 Ligand / CD154 / TRAP1 (Activation Marker of T-Lymphocytes) Antibody, IgG2b, Clone: [CD40LG/2761], Mouse, Monoclonal |
| Artikelnummer | NBT-959-MSM1-B1 |
| Hersteller Artikelnummer | 959-MSM1-B1 |
| Alternativnummer | NBT-959-MSM1-B1-0.5,NBT-959-MSM1-B1-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | ELISA, FC, IF, IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment (around aa 108-261) of human CD40LG protein (exact sequence is proprietary) |
| Produktbeschreibung | CD40LG expression is mainly confined to the CD4-positive-T-cell subset. Its expression is induced shortly after T-cell activation and represents an early activation marker of T lymphocytes. CD40 is constitutively expressed mainly on B cells, macropha... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [CD40LG/2761] |
| Molekulargewicht | 36kDa |

| | |
|------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Isotyp | IgG2b |
| NCBI | 959 |
| UniProt | P29965 |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | ELISA (For coating, order antibody without BSA), Flow Cytometry (2-4ug/million cells), Immunofluorescence (2-4ug/ml), Western Blot (2-4ug/ml), Immunohistochemistry (Formalin-fixed) (2-4ug/ml for 30 minutes at RT), (Staining of formalin-fixed tissues requ |