

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Plasma Cell Marker Antibody, IgG2a, Clone: [LIV3G11, same as 7B18], Mouse, Monoclonal NBT-MSM1-47-P0

| | |
|--------------------------|---|
| Artikelname | Plasma Cell Marker Antibody, IgG2a, Clone: [LIV3G11, same as 7B18], Mouse, Monoclonal |
| Artikelnummer | NBT-MSM1-47-P0 |
| Hersteller Artikelnummer | MSM1-47-P0 |
| Alternativnummer | NBT-MSM1-47-P0-20,NBT-MSM1-47-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Pancreatic cancer-related mucin |
| Produktbeschreibung | It recognizes an intra-cytoplasmic antigen, which shows a very high degree of specificity for plasma cells. This antigen is present in normal as well as neoplastic plasma cells. Plasma cells, which are large lymphocytes derived from an antigen-specif... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [LIV3G11, same as 7B18] |
| Molekulargewicht | Not Known |
| Isotyp | IgG2a |

| | |
|------------------------|---|
| NCBI | Not Known |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (0.1-0.2ug/ml for 30 minutes at RT) [(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilu |