

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Melanoma Marker (MART-1 + Tyrosinase + gp100) Antibody, Clone: [M2-7C10 + M2-9E3 + T311 + HMB45], Mouse, Monoclonal NBT-MSM1-701-P1ABX

Artikelname	Melanoma Marker (MART-1 + Tyrosinase + gp100) Antibody, Clone: [M2-7C10 + M2-9E3 + T311 + HMB45], Mouse, Monoclonal
Artikelnummer	NBT-MSM1-701-P1ABX
Hersteller Artikelnummer	MSM1-701-P1ABX
Alternativnummer	NBT-MSM1-701-P1ABX-100
Hersteller	NeoBiotechnologies
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant hMART-1 protein (M2-7C10, M2-9E3), Recombinant tyrosinase protein (T311), Extract of pigmented melanoma metastases from lymph nodes (HMB45)
Produktbeschreibung	This antibody cocktail recognizes three melanoma-specific proteins, which include MART-1, Tyrosinase and gp100. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Tyrosinase is one of the...
Klonalität	Monoclonal
Klon-Bezeichnung	[M2-7C10 + M2-9E3 + T311 + HMB45]

Molekulargewicht	20-22kDa (doublet) (MART), 70-80kDa (Tyrosinase), 26-100kDa (gp100)
NCBI	<a href="#">2315</a>
UniProt	<a href="#">Q16655</a>
Formulierung	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Antibody Type	Monoclonal Antibody
Anwendungsbeschreibung	Western Blot (2-4ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 min