

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Negative Control for Rabbit Monoclonal Antibodies Antibody, Clone: [NCRBM/1520R] NBT-RBM1-1520-P0

| | |
|--------------------------|--|
| Artikelname | Negative Control for Rabbit Monoclonal Antibodies Antibody, Clone: [NCRBM/1520R] |
| Artikelnummer | NBT-RBM1-1520-P0 |
| Hersteller Artikelnummer | RBM1-1520-P0 |
| Alternativnummer | NBT-RBM1-1520-P0-20,NBT-RBM1-1520-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | ELISA, FC, IF, IHC, WB |
| Spezies Reaktivität | All |
| Immunogen | Recombinant full-length human KRT7 protein |
| Produktbeschreibung | To serve as a true negative control for various procedures, this rabbit monoclonal IgG should be used under the identical conditions and at the same dilution as the rabbit monoclonal antibody being used for a positive reaction.... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [NCRBM/1520R] |
| Molekulargewicht | Not Applicable |
| NCBI | Not Applicable |

| | |
|------------------------|--|
| Formulierung | 200ug/ml of Ab purified by Protein A Column. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | ELISA, Flow Cytometry, Immunofluorescence, Immunoprecipitation, Western Blot (1-2ug/ml), Immunohistochemistry (Frozen), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue section |