

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant Bromodeoxyuridine (BrdU) (Proliferation Marker) Antibody, Clone: [BRDU/3902R], Rabbit, Monoclonal NBT-RBM13-3902-P1ABX

| | |
|--------------------------|---|
| Artikelname | Recombinant Bromodeoxyuridine (BrdU) (Proliferation Marker) Antibody, Clone: [BRDU/3902R], Rabbit, Monoclonal |
| Artikelnummer | NBT-RBM13-3902-P1ABX |
| Hersteller Artikelnummer | RBM13-3902-P1ABX |
| Alternativnummer | NBT-RBM13-3902-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | All |
| Immunogen | Bromodeoxyuridine (BrdU) conjugated to KLH |
| Produktbeschreibung | It reacts with Bromodeoxyuridine (BrdU) in single stranded DNA (produced by partial denaturation of double stranded DNA), BrdU coupled to a protein carrier, as well as free BrdU. BrdU is a thymidine analog, incorporated into cell nuclei during DNA sy... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [BRDU/3902R] |
| Molekulargewicht | Depends on the target |
| NCBI | Not Applicable |

| | |
|------------------------|--|
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT) [For staining of formalin-fixed tissues, incubate sections in 4N HCl for 30 minutes at RT followed by digestion with trypsin at 1mg/ml PBS, 10 min at 37oC] Optimal dilution for a speci |