

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

PECAM-1 Antibody [C31.7], Unconjugated, Mouse, Monoclonal PRS-33-185

Artikelname	PECAM-1 Antibody [C31.7], Unconjugated, Mouse, Monoclonal
Artikelnummer	PRS-33-185
Hersteller Artikelnummer	33-185
Alternativnummer	PRS-33-185-100
Hersteller	ProSci
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human, Primate, Rabbit
Immunogen	Human recombinant PECAM-1 / CD31 protein was used as the immunogen for this antibody.
Konjugation	Unconjugated
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[C31.7]
NCBI	5175
UniProt	P16284
Puffer	PBS with 0.1 mg/ml BSA and 0.05% sodium azide

Formulierung	Liquid
Application Verdünnung	Optimal dilutions for each application to be determined by the researcher
Anwendungsbeschreibung	<p>Flow Cytometry: 0.5-1ug/10e6 cells IF: 1-2 ug/ml WB: 0.5-1 ug/ml IHC (FFPE): 0.5-1 ug/ml for 30 minutes at RT (1) Prediluted format : incubate for 30 min at RT (2) The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the PECAM-1 antibody to be titered up or down for optimal performance.</p> <p>1. Staining of formalin-fixed tissues requires boiling tissue sections in 1mM EDTA, pH 7.5-8.5, for 10-20 min followed by cooling at RT for 20 minutes.</p> <p>2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p>