

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### CD45RA Antibody [158-4D3], Unconjugated, Mouse, Monoclonal PRS-33-195

Artikelname	CD45RA Antibody [158-4D3], Unconjugated, Mouse, Monoclonal
Artikelnummer	PRS-33-195
Hersteller Artikelnummer	33-195
Alternativnummer	PRS-33-195-100
Hersteller	ProSci
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human
Immunogen	Stimulated human leukocytes were used as the immunogen for this CD45RA antibody.
Konjugation	Unconjugated
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[158-4D3]
NCBI	<a href="#">5788</a>
UniProt	<a href="#">P08575</a>
Puffer	PBS with 0.1 mg/ml BSA and 0.05% sodium azide

Formulierung	Liquid
Application Verdünnung	Optimal dilutions for each application to be determined by the researcher
Anwendungsbeschreibung	<p>Flow Cytometry: 0.5-1 ug/million cells  IF: 1-2 ug/ml  IHC (FFPE): 0.5-1 ug/ml for 30 minutes at RT (1)  Prediluted format : incubate for 30 min at RT (2)  The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the CD45RA antibody to be titered up or down for optimal performance.</p> <p>1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes.</p> <p>2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p>