

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Histone Antibody [1415-1], Unconjugated, Mouse, Monoclonal PRS-33-387

Artikelname	Histone Antibody [1415-1], Unconjugated, Mouse, Monoclonal
Artikelnummer	PRS-33-387
Hersteller Artikelnummer	33-387
Alternativnummer	PRS-33-387-100
Hersteller	ProSci
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC-P, WB
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Nuclei of human leukemia biopsy cells were used as the immunogen for the Histone antibody.
Konjugation	Unconjugated
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[1415-1]
NCBI	<a href="#">3005</a>
UniProt	<a href="#">P07305</a>
Puffer	PBS with 0.1 mg/ml BSA and 0.05% sodium azide

Formulierung	Liquid
Application Verdünnung	Optimal dilutions for each application to be determined by the researcher
Anwendungsbeschreibung	<p>Western blot: 0.5-1 ug/ml  Flow Cytometry: 0.5-1 ug/million cells in 0.1ml  Immunofluorescence: 0.5-1 ug/ml  Immunohistochemistry (FFPE): 0.5-1 ug/ml for 30 min at RT (1)  Prediluted format : incubate for 30 min at RT (2)  Optimal dilution of the Histone antibody should be determined by the researcher.</p> <p>1. Staining of formalin/paraffin tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.</p> <p>2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p>