

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Cytokeratin 19 Antibody [KRT19/799], Unconjugated, Mouse, Monoclonal PRS-33-434

Artikelname	Cytokeratin 19 Antibody [KRT19/799], Unconjugated, Mouse, Monoclonal
Artikelnummer	PRS-33-434
Hersteller Artikelnummer	33-434
Alternativnummer	PRS-33-434-100
Hersteller	ProSci
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC-P, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human KRT19 protein was used as the immunogen for the Cytokeratin 19 antibody.
Konjugation	Unconjugated
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[KRT19/799]
NCBI	<a href="#">3880</a>
UniProt	<a href="#">P08727</a>
Puffer	PBS with 0.1 mg/ml BSA and 0.05% sodium azide

Formulierung	Liquid
Application Verdünnung	Optimal dilutions for each application to be determined by the researcher
Anwendungsbeschreibung	<p>Flow Cytometry: 0.5-1 ug/million cells in 0.1ml  Immunofluorescence: 1-2 ug/ml  Immunohistochemistry (FFPE): 0.5-1 ug/ml for 30 min at RT  (1) Prediluted format : incubate for 30 min at RT (2) Optimal dilution of the Cytokeratin 19 antibody should be determined by the researcher.</p> <p>1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min  2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p>