

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

MFGE8 Antibody [MFG-06], Unconjugated, Mouse, Monoclonal PRS-33-436

Artikelname	MFGE8 Antibody [MFG-06], Unconjugated, Mouse, Monoclonal
Artikelnummer	PRS-33-436
Hersteller Artikelnummer	33-436
Alternativnummer	PRS-33-436-100
Hersteller	ProSci
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC-P, WB
Spezies Reaktivität	Canine, Human
Immunogen	A human milk fat globule membrane preparation was used as the immunogen for the MFGE8 antibody.
Konjugation	Unconjugated
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[MFG-06]
NCBI	4240
UniProt	Q08431
Puffer	PBS with 0.1 mg/ml BSA and 0.05% sodium azide

Formulierung	Liquid
Application Verdünnung	Optimal dilutions for each application to be determined by the researcher
Anwendungsbeschreibung	<p>Western blot: 1-2 ug/ml Flow Cytometry: 0.5-1 ug/million cells in 0.1ml Immunofluorescence: 0.5-1 ug/ml Immunohistochemistry (FFPE): 0.5-1 ug/ml for 30 min at RT (1) Prediluted format : incubate for 30 min at RT (2) Optimal dilution of the MFGE8 antibody should be determined by the researcher.</p> <p>1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p>