

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Beta-2 Microglobulin Antibody [B2M/961], Unconjugated, Mouse, Monoclonal PRS-33-458

Artikelname	Beta-2 Microglobulin Antibody [B2M/961], Unconjugated, Mouse, Monoclonal
Artikelnummer	PRS-33-458
Hersteller Artikelnummer	33-458
Alternativnummer	PRS-33-458-100
Hersteller	ProSci
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC-P
Spezies Reaktivität	Human, Primate
Immunogen	Full length recombinant human protein was used as the immunogen for the Beta-2 Microglobulin antibody.
Konjugation	Unconjugated
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[B2M/961]
NCBI	567
UniProt	P61769

Puffer	PBS with 0.1 mg/ml BSA and 0.05% sodium azide
Formulierung	Liquid
Application Verdünnung	Optimal dilutions for each application to be determined by the researcher
Anwendungsbeschreibung	<p>Flow Cytometry: 0.5-1 ug/million cells in 0.1ml</p> <p>Immunofluorescence: 0.5-1 ug/ml</p> <p>Immunohistochemistry (FFPE): 0.5-1 ug/ml for 30 min at RT (1)</p> <p>Prediluted format: incubate for 30 min at RT (2)</p> <p>Optimal dilution of the Beta-2 Microglobulin antibody should be determined by the researcher.</p> <p>1. Staining of formalin/paraffin tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.</p> <p>2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p>