

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### FGF-4 Antibody (biotin), Biotin, Rabbit, Polyclonal PRS-38-122

|                          |  |
|--------------------------|--|
| Artikelname              | FGF-4 Antibody (biotin), Biotin, Rabbit, Polyclonal  |
| Artikelnummer            | PRS-38-122   |
| Hersteller Artikelnummer | 38-122   |
| Alternativnummer         | PRS-38-122-0.05  |
| Hersteller               | ProSci   |
| Wirt                     | Rabbit   |
| Kategorie                | Antikörper   |
| Applikation              | ELISA, WB  |
| Spezies Reaktivität      | Human  |
| Immunogen                | Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hFGF-4. Human FGF-4 specific antibody was purified by affinity chromatography and then biotinylated. |
| Konjugation              | Biotin   |
| Klonalität               | Polyclonal   |
| Konzentration            | batch dependent  |
| NCBI                     | <a href="#">2249</a>   |
| UniProt                  | <a href="#">P08620</a>   |
| Formulierung             | Lyophilized  |
| Application Verdünnung   | Centrifuge vial prior to opening.  |

Anwendungsbeschreibung

ELISA:Sandwich:To detect hFGF-4 by sandwich ELISA (using 100  $\mu$ L/well antibody solution) a concentration of 0.25 - 1.0  $\mu$ g/mL of this antibody is required. This biotinylated polyclonal antibody, in conjunction with our polyclonal Anti-Human FGF-4 as a capture antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hFGF-4. Western Blot:To detect hFGF-4 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2  $\mu$ g/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant hFGF-4 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.