

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cystatin C Peptide PRS-45-043P

Artikelname	Cystatin C Peptide
Artikelnummer	PRS-45-043P
Hersteller Artikelnummer	45-043P
Alternativnummer	PRS-45-043P-0.1
Hersteller	ProSci
Kategorie	Proteine/Peptide
Applikation	Blocking
NCBI	13010
UniProt	P21460
Puffer	Cystatin C peptide is supplied as 100 µg of dried peptide. When the peptide is reconstituted in 200 µL water, the concentration will be 0.5 mg/mL.
Formulierung	Lyophilized

Anwendungsbeschreibung

Cystatin C Peptide is used for blocking the activity of Cystatin C antibody (catalog number 45-043). The Cystatin C antibody ships at 0.5 mg/mL and the Cystatin C Peptide ships as a 100 ug pellet. When peptide is reconstituted in 200 uL water, the concentration will be 0.5 mg/mL, identical to that of the antibody. The best starting ratio would be 1:1 (which means molar excess of peptide relative to antibody when identical volumes are mixed). Mix equal volume of peptide and antibody at the required dilution and leave at ambient temperature for 20 minutes. Then incubate and develop in parallel two blots: one with the antibody and one with the antibody which has been pre-absorbed with the provided peptide.