

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Flotillin 2 Peptide PRS-45-052P

Artikelname	Flotillin 2 Peptide
Artikelnummer	PRS-45-052P
Hersteller Artikelnummer	45-052P
Alternativnummer	PRS-45-052P-0.1
Hersteller	ProSci
Kategorie	Proteine/Peptide
Applikation	Blocking
NCBI	2319
UniProt	Q14254
Puffer	Flotillin 2 peptide is supplied as 100 µg of dried peptide. When the peptide is reconstituted in 200 µL water, the concentration will be 0.5 mg/mL.
Formulierung	Lyophilized

Anwendungsbeschreibung

Flotillin 2 Peptide is used for blocking the activity of Flotillin 2 antibody (catalog number 45-052). The Flotillin 2 antibody ships at 0.5 mg/mL and the Flotillin 2 Peptide ships as a 100 ug pellet. When peptide is reconstituted in 200 uL water, the concentration will be 0.5 mg/mL, identical to that of the antibody. The best starting ratio would be 1:1 (which means molar excess of peptide relative to antibody when identical volumes are mixed). Mix equal volume of peptide and antibody at the required dilution and leave at ambient temperature for 20 minutes. Then incubate and develop in parallel two blots: one with the antibody and one with the antibody which has been pre-absorbed with the provided peptide.