

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### MTA1 Peptide PRS-45-094P

|                          |                                                                                                                                              |
|--------------------------|----------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | MTA1 Peptide                                                                                                                                 |
| Artikelnummer            | PRS-45-094P                                                                                                                                  |
| Hersteller Artikelnummer | 45-094P                                                                                                                                      |
| Alternativnummer         | PRS-45-094P-0.1                                                                                                                              |
| Hersteller               | ProSci                                                                                                                                       |
| Kategorie                | Proteine/Peptide                                                                                                                             |
| Applikation              | Blocking                                                                                                                                     |
| NCBI                     | <a href="#">9112</a>                                                                                                                         |
| UniProt                  | <a href="#">Q13330</a>                                                                                                                       |
| Puffer                   | MTA1 peptide is supplied as 100 µg of dried peptide. When the peptide is reconstituted in 200 µL water, the concentration will be 0.5 mg/mL. |
| Formulierung             | Lyophilized                                                                                                                                  |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Anwendungsbeschreibung | <p>MTA1 Peptide is used for blocking the activity of MTA1 antibody (catalog number 45-094). The MTA1 antibody ships at 0.5 mg/mL and the MTA1 Peptide ships as a 100 ug pellet. When peptide is reconstituted in 200 uL water, the concentration will be 0.5 mg/mL, identical to that of the antibody. The best starting ratio would be 1:1 (which means molar excess of peptide relative to antibody when identical volumes are mixed). Mix equal volume of peptide and antibody at the required dilution and leave at ambient temperature for 20 minutes. Then incubate and develop in parallel two blots: one with the antibody and one with the antibody which has been pre-absorbed with the provided peptide.</p> |
|------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|