

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

PDL2 Antibody [7C7], IgG1, Unconjugated, Mouse, Monoclonal PRS-RF16023

| | |
|--------------------------|--|
| Artikelname | PDL2 Antibody [7C7], IgG1, Unconjugated, Mouse, Monoclonal |
| Artikelnummer | PRS-RF16023 |
| Hersteller Artikelnummer | RF16023 |
| Alternativnummer | PRS-RF16023-0.02, PRS-RF16023-0.1 |
| Hersteller | ProSci |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | ELISA, FC, ICC, IF, IHC-P, WB |
| Spezies Reaktivität | Human |
| Immunogen | PD-L2 antibody was raised against the extracellular domain of human PD-L2. |
| Konjugation | Unconjugated |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/mL |
| Klon-Bezeichnung | [7C7] |
| Molekulargewicht | Predicted: 30 kDa Observed: 38 kDa |
| Isotyp | IgG1 |
| NCBI | 80380 |

| | |
|------------------------|---|
| UniProt | P9BQ51 |
| Puffer | PD-L2 Antibody is supplied in PBS containing 0.02% sodium azide and 50% glycerol. |
| Formulierung | Liquid |
| Application Verdünnung | Optimal dilutions for each application to be determined by the researcher. |
| Anwendungsbeschreibung | PD-L2 antibody can be used for detection of PD-L2 by Western blot at 0.5 - 1 µg/mL. Antibody can also be used for immunohistochemistry starting at 2 µg/mL. For immunofluorescence start at 20 µg/mL. Antibody validated: Western Blot in human samples, Immunohistochemistry in human samples, Immunocytochemistry in human samples, Immunofluorescence in human samples and Flow Cytometry in mouse samples. All other applications and species not yet tested. |