

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Eotaxin-2 Antibody, Unconjugated, Goat, Polyclonal PRS-XP-5125

Artikelname	Eotaxin-2 Antibody, Unconjugated, Goat, Polyclonal
Artikelnummer	PRS-XP-5125
Hersteller Artikelnummer	XP-5125
Alternativnummer	PRS-XP-5125-0.1
Hersteller	ProSci
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA, NeA, WB
Spezies Reaktivität	Mouse
Immunogen	Produced from sera of goats pre-immunized with highly pure (>98%) recombinant murine Eotaxin-2.
Konjugation	Unconjugated
Klonalität	Polyclonal
Konzentration	batch dependent
NCBI	56221
UniProt	Q9JKC0
Formulierung	Lyophilized
Application Verdünnung	Centrifuge vial prior to opening.

Anwendungsbeschreibung

Neutralization: To yield one-half maximal inhibition [ND50] of the biological activity of mEotaxin-2 (100 ng/mL), a concentration of 7.0-8.0 µg/mL of this antibody is required. ELISA: To detect murine Eotaxin-2 by direct ELISA (using 100 µL/well antibody solution) a concentration of at least 0.5 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of 0.2 - 0.4 ng/well of recombinant murine Eotaxin-2. Sandwich: To detect mEotaxin-2 by sandwich ELISA (using 100 µL/well antibody solution) a concentration of 0.5 - 2.0 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with our Biotinylated Anti-Murine Eotaxin-2 (XP-5125Bt) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant mEotaxin-2. Western Blot: To detect murine Eotaxin-2 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant murine Eotaxin-2 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.