

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### MIG Antibody, Unconjugated, Rabbit, Polyclonal PRS-XP-5231

Artikelname	MIG Antibody, Unconjugated, Rabbit, Polyclonal
Artikelnummer	PRS-XP-5231
Hersteller Artikelnummer	XP-5231
Alternativnummer	PRS-XP-5231-0.1
Hersteller	ProSci
Wirt	Rabbit
Kategorie	Antikörper
Applikation	ELISA, NeA, WB
Spezies Reaktivität	Human
Immunogen	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hMIG (human MIG).
Konjugation	Unconjugated
Klonalität	Polyclonal
Konzentration	batch dependent
NCBI	<a href="#">4283</a>
UniProt	<a href="#">Q07325</a>
Formulierung	Lyophilized
Application Verdünnung	Centrifuge vial prior to opening.

Anwendungsbeschreibung

Neutralization: To yield one-half maximal inhibition [ND50] of the biological activity of hMIG (100.00 ng/mL), a concentration of 5.0 - 7.0 µg/mL of this antibody is required. ELISA: To detect hMIG by direct ELISA (using 100 µL/well antibody solution) a concentration of at least 0.5 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of 0.2 - 0.4 ng/well of recombinant hMIG. Sandwich: To detect hMIG by sandwich ELISA (using 100 µL/well antibody solution) a concentration of 0.5 - 2.0 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with our Biotinylated Anti-Human MIG (XP-5231Bt) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hMIG. Western Blot: To detect hMIG by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant hMIG is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.