

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD56 / NCAM(123C3.D5), CF405S conjugate, 0.1mg/mL, Clone: [123C3.D5], Mouse, Monoclonal BOT-BNC040015-100**

Artikelname	CD56 / NCAM(123C3.D5), CF405S conjugate, 0.1mg/mL, Clone: [123C3.D5], Mouse, Monoclonal
Artikelnummer	BOT-BNC040015-100
Hersteller Artikelnummer	BNC040015-100
Alternativnummer	BOT-BNC040015-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC
Spezies Reaktivität	Human
Immunogen	Membrane preparation of a small cell lung carcinoma
Konjugation	CF405S
Produktbeschreibung	This MAb reacts with an extracellular domain (close to transmembrane) of CD56/NCAM. Three isoforms of neural cell adhesion molecule (NCAM) are produced by differential splicing of the RNA transcript from a single gene. The 135 kDa isoform is the basi...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[123C3.D5]
Molekulargewicht	180, 145 and 125 kDa
UniProt	<a href="#">P13591</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user