

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Ep-CAM / CD326(VU-1D9), CF405S conjugate, 0.1mg/mL, Clone: [VU-1D9], Mouse, Monoclonal BOT-BNC040017-100

| | |
|--------------------------|---|
| Artikelname | Ep-CAM / CD326(VU-1D9), CF405S conjugate, 0.1mg/mL, Clone: [VU-1D9], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC040017-100 |
| Hersteller Artikelnummer | BNC040017-100 |
| Alternativnummer | BOT-BNC040017-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | Exosome Staining, FC, IF, IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Small cell lung carcinoma cells |
| Konjugation | CF405S |
| Produktbeschreibung | This antibody reacts with the first EGF repeat in the extracellular domain of Ep-CAM. It is a 40-43 kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [VU-1D9] |

| | |
|------------------------|---|
| Molekulargewicht | 40-43 kDa |
| UniProt | P16422 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Does not react with rat or ferret, others not known Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |