

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD50 (ICAM-3)(186-2G9), CF405S conjugate, 0.1mg/mL, Clone: [186-2G9], Mouse, Monoclonal BOT-BNC040178-100**

Artikelname	CD50 (ICAM-3)(186-2G9), CF405S conjugate, 0.1mg/mL, Clone: [186-2G9], Mouse, Monoclonal
Artikelnummer	BOT-BNC040178-100
Hersteller Artikelnummer	BNC040178-100
Alternativnummer	BOT-BNC040178-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IF, IHC
Spezies Reaktivität	Human
Immunogen	Stimulated human leukocytes
Konjugation	CF405S
Produktbeschreibung	Recognizes the D1 domain of an N-glycosylated glycoprotein of 120 kDa with intra-chain disulfide bonds, identified as CD50 or ICAM-3 (Leucocyte Workshop VI). CD50 is the major ligand for LFA-1 (CD11a/CD18) and may have signaling role to increase adhe...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[186-2G9]

Molekulargewicht	110-160 kDa
UniProt	<a href="#">P32942</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Blocks binding of CD11a LFA-1 to CD50 ICAM-3 Optimal dilution for a specific application should be determined by user