

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **FOXP3(FXP3/197), CF405S conjugate, 0.1mg/mL, Clone: [FXP3/197], Mouse, Monoclonal BOT-BNC040197-500**

|                          |   |
|--------------------------|---|
| Artikelname              | FOXP3(FXP3/197), CF405S conjugate, 0.1mg/mL, Clone: [FXP3/197],<br>Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNC040197-500   |
| Hersteller Artikelnummer | BNC040197-500   |
| Alternativnummer         | BOT-BNC040197-500-500UL   |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human, Monkey, Mouse  |
| Immunogen                | Recombinant human full-length FOXP3 protein   |
| Konjugation              | CF405S  |
| Produktbeschreibung      | Recognizes a protein of 47-55 kDa, which is identified as FOXP3. Its precise epitope is not known, but it has been mapped to the N-terminal portion of the protein. The FOX family of transcription factors is a large group of proteins that share a comm... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [FXP3/197]  |

|                        |  |
|------------------------|--|
| Molekulargewicht       | 47-55 kDa  |
| UniProt                | <a href="#">Q9BZS1</a>   |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide   |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris buffer with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |