

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Complement C4d(C4D204), CF405S conjugate, 0.1mg/mL, Clone: [C4D204], Mouse, Monoclonal BOT-BNC040204-100

| | |
|--------------------------|---|
| Artikelname | Complement C4d(C4D204), CF405S conjugate, 0.1mg/mL, Clone: [C4D204], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC040204-100 |
| Hersteller Artikelnummer | BNC040204-100 |
| Alternativnummer | BOT-BNC040204-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IF, IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human Complement 4d protein |
| Konjugation | CF405S |
| Produktbeschreibung | This MAb is specific to Complement 4d (C4d) and it reacts with the secreted as well as cell-bound C4d. C4d is a degradation product of the activated complement factor C4b. Complement 4b is typically activated by binding of Abs to specific target mole... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [C4D204] |

| | |
|------------------------|--|
| Molekulargewicht | 192 kDa (predicted) |
| UniProt | P0COL4 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology formalin-fixed 1:200-1:400 Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Immunofluorescence 1:50-1:100 ELISA For coating use Ab at 1-2 ug/mL, order Ab without BSA Optimal dilution for a specific application should be determined by user |