

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD8A (Cytotoxic / Suppressor T-Cell Marker)(T8/209), CF405S conjugate, 0.1mg/mL, Clone: [T8/209], Mouse, Monoclonal BOT-BNC040209-500**

Artikelname	CD8A (Cytotoxic / Suppressor T-Cell Marker)(T8/209), CF405S conjugate, 0.1mg/mL, Clone: [T8/209], Mouse, Monoclonal
Artikelnummer	BOT-BNC040209-500
Hersteller Artikelnummer	BNC040209-500
Alternativnummer	BOT-BNC040209-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Spezies Reaktivität	Human, Primate
Immunogen	Recombinant human T8 (CD8a) protein
Konjugation	CF405S
Produktbeschreibung	This MAb is specific for the 32 kDa alpha chain of human CD8, known as CD8a or CD8 alpha. CD8 is a cell surface receptor expressed either as a heterodimer with the CD8 beta chain (CD8 alpha/beta) or as a homodimer (CD8 alpha/alpha). A majority of thy...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[T8/209]
Molekulargewicht	32 kDa

UniProt	<a href="#">P01732</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry