

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cytokeratin, pan(AE-1 / AE-3), CF405S conjugate, 0.1mg/mL, Clone: [AE-1 / AE-3], Mouse, Monoclonal BOT-BNC040371-100**

Artikelname	Cytokeratin, pan(AE-1 / AE-3), CF405S conjugate, 0.1mg/mL, Clone: [AE-1 / AE-3], Mouse, Monoclonal
Artikelnummer	BOT-BNC040371-100
Hersteller Artikelnummer	BNC040371-100
Alternativnummer	BOT-BNC040371-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IHC
Spezies Reaktivität	Bovine, Canine, Gallus, Human, Monkey, Mouse, Rabbit, Rat
Immunogen	Human epidermal keratin
Konjugation	CF405S
Produktbeschreibung	Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI 6.0) subfamilies. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, which include 67 kDa (CK1), 64 kDa (CK3), ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[AE-1 / AE-3]

Molekulargewicht	40-67 kDa
UniProt	<a href="#">Q7Z794</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user