

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin, pan(AE-1 / AE-3), CF405S conjugate, 0.1mg/mL, Clone: [AE-1 / AE-3], Mouse, Monoclonal BOT-BNC040371-500

Artikelname	Cytokeratin, pan(AE-1 / AE-3), CF405S conjugate, 0.1mg/mL, Clone: [AE-1 / AE-3], Mouse, Monoclonal
Artikelnummer	BOT-BNC040371-500
Hersteller Artikelnummer	BNC040371-500
Alternativnummer	BOT-BNC040371-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IHC
Spezies Reaktivität	Bovine, Canine, Gallus, Human, Monkey, Mouse, Rabbit, Rat
Immunogen	Human epidermal keratin
Konjugation	CF405S
Produktbeschreibung	Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI 6.0) subfamilies. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, which include 67 kDa (CK1), 64 kDa (CK3), ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[AE-1 / AE-3]

Molekulargewicht	40-67 kDa
UniProt	Q7Z794
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	<p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody</p> <p>Immunofluorescence: 1-2 ug/mL</p> <p>Immunohistology formalin-fixed 0.25-0.5 ug/mL</p> <p>Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes</p> <p>Flow Cytometry 0.5-1 ug/million cells/0.1 mL</p> <p>Western blotting 0.5-1 ug/mL</p> <p>Optimal dilution for a specific application should be determined by user</p>