

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Muscle Specific Actin(HHF35), CF405S conjugate, 0.1mg/mL, Clone: [HHF35], Mouse, Monoclonal BOT-BNC040445-500

| | |
|--------------------------|---|
| Artikelname | Muscle Specific Actin(HHF35), CF405S conjugate, 0.1mg/mL, Clone: [HHF35], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC040445-500 |
| Hersteller Artikelnummer | BNC040445-500 |
| Alternativnummer | BOT-BNC040445-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Canine, Feline, Gallus, Human, Mouse, Rabbit, Rat |
| Immunogen | SDS extract of human myocardium. |
| Konjugation | CF405S |
| Produktbeschreibung | This antibody recognizes actin of skeletal, cardiac, and smooth muscle cells. It is not reactive with other mesenchymal cells except for myoepithelium. Actin can be resolved on the basis of its isoelectric points into three distinctive components: al... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [HHF35] |

| | |
|------------------------|---|
| Molekulargewicht | Multiple |
| UniProt | P62736 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |