

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin, HMW(34BE12), CF405S conjugate, 0.1mg/mL, Clone: [34BE12], Mouse, Monoclonal BOT-BNC040446-500

| | |
|--------------------------|---|
| Artikelname | Cytokeratin, HMW(34BE12), CF405S conjugate, 0.1mg/mL, Clone: [34BE12], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC040446-500 |
| Hersteller Artikelnummer | BNC040446-500 |
| Alternativnummer | BOT-BNC040446-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IF, IHC, WB |
| Spezies Reaktivität | Human, Mouse, Rat |
| Immunogen | Solubilized keratin extract from human stratum corneum |
| Konjugation | CF405S |
| Produktbeschreibung | This antibody recognizes CK1, CK5, CK10 and CK14. In normal epithelia, it stains stratified epithelia, myoepithelial cells and basal cells in the prostate gland and bronchi. This MAb shows no reactivity with hepatocytes, pancreatic acinar cells, prox... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [34BE12] |

| | |
|------------------------|---|
| Molekulargewicht | 50-67 kDa |
| UniProt | Q01546 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Immunofluorescence 0.5-1.0 ug/mL Western blotting 0.5-1.0 ug/mL Optimal dilution for a specific application should be determined by user |